



Dietary Hyperglycemia and Its Effects on Microbial Dysbiosis and Aging in *Drosophila melanogaster*

Sunita Devi¹

Research Scholar, Department of Zoology, Shri Jagadishprasad, Jhabarmal Tibrewala University, Jhunjhunu, Rajasthan, India

Dr. Imran Khan²

Assistant Professor, Department of Zoology, Shri Jagadishprashad, Jhabarmal Tibrewala University, Jhunjhunu, Rajasthan, India

Abstract:

This study examines the effects of dietary hyperglycemia on gut microbiota dysbiosis and aging in *Drosophila melanogaster*. Flies were maintained on control and high-sugar diets and analyzed across different age stages. Results revealed a significant reduction in microbial diversity under high-sugar conditions, with depletion of beneficial bacteria such as *Lactobacillus* and increased abundance of opportunistic taxa including *Enterobacter* and *Serratia*. Elevated bacterial load and pronounced dysbiosis were observed, particularly in older flies. Physiological analysis showed reduced body size, impaired metabolic efficiency, and accelerated somatic decline. Reproductive parameters, including fecundity and offspring viability, were significantly decreased in hyperglycemic groups. Statistical analyses confirmed that both diet and age, along with their interaction, strongly influence microbial composition and host fitness. Overall, the findings highlight that high-sugar diets disrupt gut microbial balance and accelerate aging-related physiological decline, emphasizing the importance of diet in regulating host-microbiota interactions.

Keywords: Hyperglycemia, Gut microbiota, Dysbiosis, Aging, *Drosophila melanogaster*, High-sugar diet

Introduction

The growing prevalence of high-sugar diets in modern societies has raised significant concerns regarding their impact on metabolic health and overall physiological functioning. Excessive intake of dietary sugars, particularly glucose and fructose, is strongly associated with the development of hyperglycemia, obesity, insulin resistance, and other metabolic disorders. Beyond systemic metabolic effects, emerging evidence suggests that dietary sugar plays a crucial role in shaping the composition and function of the gut microbiota, a complex community of microorganisms that reside within the gastrointestinal tract. The gut microbiome is now recognized as a key regulator of host physiology, influencing nutrient metabolism, immune responses, and even reproductive performance. Therefore, understanding the interplay between dietary hyperglycemia, gut microbiota, and host aging has become a central focus in contemporary biological research (Wong et al., 2014; Baenas & Wagner, 2022).



Drosophila melanogaster has emerged as a powerful and widely used model organism for studying diet–microbiota interactions and metabolic disorders. The species offers several advantages, including a short life cycle, ease of maintenance, well-characterized genetics, and conserved metabolic pathways that are comparable to those in mammals. Importantly, the gut microbiota of *Drosophila* is relatively simple, typically dominated by a few bacterial genera such as *Lactobacillus* and *Acetobacter*, making it an ideal system for investigating microbial dynamics under controlled experimental conditions (Broderick & Lemaitre, 2012; Ridley et al., 2012). Previous studies have demonstrated that dietary composition directly influences microbial community structure in *Drosophila*, thereby affecting host development, metabolism, and lifespan. For instance, alterations in macronutrient balance can lead to shifts in microbial diversity, which in turn influence host physiological outcomes (Henry et al., 2020).

Figure 1 illustrates metabolic organs relevant to the use of *Drosophila* in studies of diet and aging (Gáliková & Klepsatel, 2018).




Figure 1. Metabolic organs in *Drosophila* and their functional human counterparts
Gáliková and Klepsatel (2018)

metabolic dysfunction in *Drosophila melanogaster*, including phenotypes analogous to obesity and type 2 diabetes in humans. Experimental studies have reported that flies fed with sugar-rich diets exhibit elevated hemolymph glucose levels, increased lipid accumulation, and impaired insulin signaling pathways (Palanker Musselman et al., 2011; van Dam et al., 2020). These



metabolic disturbances are often accompanied by significant changes in the gut microbiota, suggesting a close relationship between dietary sugar and microbial composition. In particular, hyperglycemic conditions tend to reduce microbial diversity and promote the growth of opportunistic bacteria, leading to a state of dysbiosis. Such dysbiotic conditions are associated with impaired gut function, increased oxidative stress, and reduced host fitness (Clark et al., 2015; Zhang et al., 2017).

The fat-body images in Figure 2 provide anatomical context for energy storage and metabolic dysfunction (Gáliková & Klepsatel, 2018).




Figure 2. Drosophila fat body under bright-field and confocal microscopy

The gut microbiota functions as a dynamic ecological system, where microbial species interact through cooperative and competitive relationships. Beneficial bacteria such as *Lactobacillus plantarum* and *Acetobacter pomorum* play important roles in maintaining metabolic homeostasis by facilitating nutrient absorption, producing essential metabolites, and regulating hormonal pathways involved in growth and reproduction (Erkosar et al., 2013; Consuegra et al., 2020). However, high-sugar diets can disrupt these interactions by altering the availability of nutrients and changing the gut environment. Excess sugar creates conditions that favor the proliferation of fast-growing, sugar-metabolizing bacteria, often at the expense of beneficial commensals. This shift from a balanced microbial community to a dysbiotic state can impair nutrient utilization and reduce the efficiency of metabolic processes (Whon et al., 2017).

In addition to metabolic effects, dietary hyperglycemia has been shown to influence reproductive fitness, which is a critical determinant of evolutionary success. Reproductive traits such as fecundity, ovariole number, and offspring viability are highly sensitive to nutritional status and energy availability. In *Drosophila*, optimal reproductive performance depends on efficient nutrient allocation and proper hormonal regulation. Studies have demonstrated that excessive sugar intake can negatively affect reproductive output by disrupting energy balance and altering signaling pathways involved in oogenesis (Lushchak et al., 2014; Klepsatel et al., 2020). Furthermore,



microbial dysbiosis induced by high-sugar diets may indirectly affect reproduction by impairing metabolic efficiency and increasing physiological stress.

Aging is another important factor that influences both gut microbiota composition and host physiology. As organisms age, natural changes occur in microbial diversity and community structure, often resulting in reduced stability and increased susceptibility to dysbiosis. In *Drosophila melanogaster*, aging has been associated with shifts in gut microbiota that impair intestinal function and contribute to decreased lifespan (Clark et al., 2015). When combined with dietary stress, such as high-sugar intake, these age-related changes may be amplified, leading to accelerated physiological decline. Recent research suggests that diet and age interact synergistically to influence microbial composition, with high-sugar diets intensifying age-related dysbiosis and metabolic dysfunction (Arias-Rojas & Iatsenko, 2022; Jelena et al., 2025).

The relationship between gut microbiota and host physiology is further mediated by microbial metabolites, which play a crucial role in regulating metabolic and reproductive processes. Microbial-derived compounds such as short-chain fatty acids, vitamins, and amino acids contribute to energy balance and influence signaling pathways such as insulin and TOR pathways (Ankrah et al., 2021). Disruption of microbial composition under hyperglycemic conditions may therefore impair these metabolic interactions, leading to reduced efficiency of nutrient utilization and compromised physiological performance. Additionally, the increase in opportunistic bacteria under high-sugar conditions may trigger inflammatory responses and oxidative stress, further exacerbating the negative effects on host health (Zhang et al., 2017).

Despite extensive research on diet-induced metabolic disorders, there remains a need for integrated studies that examine the combined effects of hyperglycemia on gut microbiota, aging, and reproductive fitness. Many previous studies have focused on individual aspects, such as microbial composition or metabolic outcomes, without considering their interconnected nature. The present study addresses this gap by adopting a comprehensive approach that simultaneously evaluates microbial diversity, physiological traits, and reproductive performance across different age stages in *Drosophila melanogaster*. By analyzing both independent and interactive effects of diet and age, the study aims to provide a deeper understanding of the mechanisms through which dietary hyperglycemia influences host-microbe interactions.

3: Research Methodology

The study is designed to investigate how high-sugar diet-induced hyperglycemia influences gut microbiota composition, physiological traits, and reproductive fitness across different age stages. A controlled laboratory experimental approach was employed to ensure accuracy, reproducibility, and validity of results.

3.2 Research Design

The study follows an experimental, comparative, and longitudinal research design. Two dietary groups—control and high-sugar diet (HSD)—were established, and observations were recorded at multiple time points (7, 21, and 45 days). This design allows for the analysis of:



- **Independent effects:** Diet and age
- **Interaction effects:** Diet $\times$ Age
- **Temporal trends:** Age-related changes under both dietary conditions

3.3 Study Organism

The model organism used in this study is *Drosophila melanogaster*, chosen for its:

- Short life cycle and ease of maintenance
- Well-characterized genome
- Conserved metabolic and aging pathways
- Simplicity of gut microbiota

Female flies were primarily used for reproductive and morphometric analyses due to their relevance in fecundity and ovarian studies.

3.4 Experimental Groups and Diet Composition

Two experimental groups were established:

1. **Control Group:**
 - Diet: Standard cornmeal–yeast–agar medium
 - Sugar concentration: ~6% w/v
 - Purpose: Baseline physiological condition
2. **High-Sugar Diet Group (HSD):**
 - Diet: Modified semi-defined medium
 - Sugar concentration: ~32% w/v (mixture of glucose and fructose)
 - Purpose: Induction of hyperglycemia and metabolic stress

Flies were maintained on these diets continuously throughout the experimental period.

4. Results and Data Analysis

The results provide insights into microbial, physiological, and reproductive responses under high-sugar dietary conditions. Each table is followed by a concise, paragraph-wise discussion.

4.1 Alpha Diversity of Gut Microbiota

Table 1: Microbial Diversity Indices under Dietary Treatments

Age (Days)	Diet	Shannon Index (H')	Simpson Index (D)	Chao1 Richness	Observed OTUs $\pm$ SD
7	Control	2.62 $\pm$ 0.07	0.85 $\pm$ 0.02	42 $\pm$ 3	45 $\pm$ 3
7	HSD	2.05 $\pm$ 0.05	0.77 $\pm$ 0.03	31 $\pm$ 2	33 $\pm$ 3
21	Control	2.48 $\pm$ 0.06	0.83 $\pm$ 0.02	40 $\pm$ 3	42 $\pm$ 2
21	HSD	1.72 $\pm$ 0.08	0.68 $\pm$ 0.04	27 $\pm$ 3	29 $\pm$ 2
45	Control	2.19 $\pm$ 0.09	0.79 $\pm$ 0.03	36 $\pm$ 2	38 $\pm$ 3
45	HSD	1.41 $\pm$ 0.07	0.58 $\pm$ 0.05	22 $\pm$ 2	25 $\pm$ 2

The diversity indices reveal a consistent and progressive decline in microbial diversity in HSD-fed flies compared to controls. At all time points, both Shannon and Simpson indices are lower in HSD groups, indicating reduced microbial richness and evenness. The effect becomes more



pronounced at later stages (45 days), suggesting that prolonged exposure to high sugar exacerbates microbial dysbiosis. The reduction in Chao1 richness and OTUs further confirms the loss of microbial complexity. In contrast, control groups maintain relatively stable diversity with only mild age-related decline. These findings indicate that dietary hyperglycemia is a dominant factor driving microbial simplification and ecosystem instability.

4.2 Beta Diversity and Community Structure

Table 2: PERMANOVA Analysis of Microbial Community Variation

Source	df	Sum of Squares	Mean Square	F-value	p-value
Diet	1	2.641	2.641	6.72	0.0008
Age	2	3.115	1.558	4.28	0.001
Diet × Age	2	1.892	0.946	3.45	0.018
Residual	12	4.712	0.392	—	—

The PERMANOVA results demonstrate that both diet and age significantly influence gut microbial community structure. The highly significant diet effect ($p = 0.0008$) indicates that sugar concentration is the primary driver of variation. Age also contributes significantly, reflecting natural temporal changes in microbial composition. The interaction term further suggests that the effect of diet intensifies with aging, resulting in increasingly distinct microbial communities in older HSD groups. These results highlight the dynamic interplay between environmental (dietary) and biological (age) factors.

4.3 Taxonomic Shifts at Phylum Level

Table 3: Relative Abundance of Major Bacterial Phyla (%)

Phylum	Control (7d)	Control (21d)	Control (45d)	HSD (7d)	HSD (21d)	HSD (45d)
Firmicutes	44.5	42.2	40.1	37.6	29.5	22.8
Proteobacteria	29.8	32.5	34.0	39.7	45.9	51.2
Bacteroidetes	18.3	17.1	15.9	14.9	12.8	10.2
Actinobacteria	7.4	8.2	10.0	7.8	11.8	15.8

A clear compositional shift is observed under high-sugar conditions, with a decrease in Firmicutes and a corresponding increase in Proteobacteria. The rise of Proteobacteria in HSD groups suggests proliferation of opportunistic bacteria associated with dysbiosis. Actinobacteria also increase with age in HSD groups, indicating adaptive microbial responses to altered gut conditions. In contrast, control groups show relatively stable phylum distribution. These findings confirm that dietary sugar induces major taxonomic restructuring.

4.4 Bacterial Load Variation

Table 4: Total Bacterial Load (CFU/ml)

Group	Mean CFU/ml ($\pm$ SD)
Control	$3.1 \times 10^5 \pm 0.4$



High-Sugar Diet	$5.6 \times 10^5 \pm 0.6$
-----------------	---------------------------

The bacterial load is significantly higher in HSD-fed flies, indicating enhanced microbial proliferation under hyperglycemic conditions. However, this increase likely reflects the growth of opportunistic taxa rather than beneficial microbes. The higher standard deviation in HSD groups suggests instability and variability in microbial populations. This pattern is indicative of dysbiotic conditions rather than healthy microbial expansion.

4.5 Reproductive Output

Table 5: Fecundity (Eggs per Female)

Age (Days)	Control	HSD
7	72 ± 6	60 ± 5
15	63 ± 5	47 ± 4
30	49 ± 4	31 ± 3

Fecundity declines with age in both groups, but the reduction is significantly greater in HSD-fed flies. The lower egg production suggests impaired ovarian function and reduced reproductive capacity. The sharper decline in HSD groups indicates accelerated reproductive aging, likely due to metabolic stress and microbial imbalance.

4.6 Developmental Viability

Table 6: Egg-to-Adult Survival (%)

Group	Viability (%)
Control	91 ± 2
High-Sugar Diet	68 ± 5

The survival rate is significantly reduced in HSD-fed flies, indicating that hyperglycemia negatively affects developmental processes. The lower viability suggests impaired embryonic development and larval growth, possibly due to metabolic stress and altered microbial interactions.

4.7 Morphometric Traits

Table 7: Thorax Length (mm)

Age	Control	HSD
7	1.18 ± 0.04	1.03 ± 0.05
21	1.14 ± 0.06	0.98 ± 0.06
45	1.05 ± 0.07	0.90 ± 0.08

Thorax length decreases with age in both groups, but the reduction is more pronounced in HSD-fed flies. This indicates impaired somatic growth under hyperglycemic conditions. The results suggest reduced anabolic activity and nutrient assimilation efficiency.

4.8 Body Weight and Size Index

Table 8: Body Weight and BSI

Age	Diet	Body Weight (mg)	BSI (mm ³)
7	Control	1.30	1.48



7	HSD	1.36	1.02
45	Control	1.12	1.08
45	HSD	0.92	0.72

HSD-fed flies show higher early body weight but lower BSI, indicating fat accumulation rather than structural growth. With age, both weight and BSI decline sharply, suggesting metabolic exhaustion. This biphasic trend reflects early overnutrition followed by physiological decline.

4.9 Reproductive Morphology

Table 9: Ovariole Number

Age	Control	HSD
7	19.2 ± 1.3	16.5 ± 1.4
21	17.1 ± 1.2	13.8 ± 1.3
45	15.3 ± 1.5	11.2 ± 1.6

Ovariole number declines significantly under HSD conditions, indicating impaired reproductive development. The reduction is more severe with age, suggesting accelerated reproductive senescence under hyperglycemia.

The results collectively demonstrate that dietary hyperglycemia significantly alters gut microbiota composition, increases bacterial load, reduces diversity, and negatively impacts physiological and reproductive traits in *Drosophila melanogaster*. The combined effects of diet and aging lead to progressive dysbiosis, metabolic imbalance, and reduced fitness, confirming the central role of diet in host–microbe interactions.

5. Conclusion

the present study demonstrates that dietary hyperglycemia induced by a high-sugar diet significantly disrupts gut microbiota composition and accelerates physiological and reproductive aging in *Drosophila melanogaster*. The findings reveal a marked decline in microbial diversity, with a shift from beneficial commensal bacteria to opportunistic taxa, indicating pronounced dysbiosis under hyperglycemic conditions. This microbial imbalance is closely associated with impaired metabolic regulation, reduced somatic growth, and a substantial decline in reproductive fitness, including decreased fecundity, ovariole number, and offspring viability. Furthermore, the interaction between diet and age intensifies these effects, leading to accelerated deterioration of both microbial stability and host physiology over time. The study highlights the critical role of diet as a primary environmental driver influencing host–microbe interactions and underscores the importance of maintaining microbial homeostasis for optimal metabolic and reproductive health. Overall, the results provide strong evidence that excessive sugar intake has detrimental long-term consequences on biological systems, emphasizing the need for balanced nutrition to preserve health and longevity.

REFERENCES

1. Huang, J. H., et al. (2015). Consumption of dietary sugar by gut bacteria determines metabolic response in *Drosophila melanogaster*. *Biology Letters*, 11(9), 20150469.



2. Zhang, X., Jin, Q., & Jin, L. H. (2017). High sugar diet disrupts gut homeostasis through JNK and STAT pathways in *Drosophila melanogaster*. *Biochemical and Biophysical Research Communications*, 487(4), 910–916.
3. Yu, C., et al. (2026). High-sugar diet induces gut microbiota dysbiosis and metabolic disruption in *Drosophila melanogaster*. *Cell Reports*.
4. Meng, Q., et al. (2024). High-sugar diet reduces intestinal microbiota and alters metabolic homeostasis in *Drosophila melanogaster*. *Process Biochemistry*.
5. Voulgari-Kokota, A., et al. (2024). High-sugar diet leads to reduced microbial diversity and impaired development in *Drosophila melanogaster*. *The ISME Journal*, 18(1).
6. Cho, K. H., et al. (2025). The gut microbiota of *Drosophila melanogaster*: A model for host–microbe interaction studies. *Microorganisms*, 13(11), 2515.
7. Zhang, Y., et al. (2025). Microbiome interactions and metabolic rhythms under high-sugar diet in *Drosophila melanogaster*. *eLife*.
8. Darby, A. M., et al. (2024). High-sugar diets increase susceptibility to bacterial infection in *Drosophila melanogaster*. *PLoS Pathogens*, 20(8), e1012447.
9. Moon, S. J., et al. (2026). Modeling tissue-specific metabolism in *Drosophila melanogaster* under hyperglycemic conditions. *Nature Communications*.
10. Qush, A., et al. (2023). Sugar-rich diet remodels gut microbiota and affects immunity and lifespan in *Drosophila melanogaster*. *Microbes and Infection*.
11. Jelena, R., et al. (2025). Long-term nutritional regimens influence gut microbiota and metabolic activity in *Drosophila melanogaster*. *Insects*, 16(2), 141.
12. Yoshinari, Y., et al. (2024). Gut hormone signaling and metabolic regulation in response to dietary sugar in *Drosophila melanogaster*. *Nature Communications*, 15(1), 10819.
13. van Dam, E., et al. (2020). Sugar-induced metabolic dysfunction and lifespan alteration in *Drosophila melanogaster*. *Cell Metabolism*, 32(2), 1–12.
14. Bombin, A., et al. (2020). Influence of diet and microbiota on metabolic phenotype in *Drosophila melanogaster*. *Microorganisms*, 8(12), 1972.
15. Guo, Y., et al. (2021). High sugar diet alters gut microbiota and metabolic pathways. *Translational Psychiatry*, 11, 267.
16. Wang, M., et al. (2022). High-sugar diet induces hyperglycemia and metabolic syndrome in *Drosophila melanogaster*. *Frontiers in Pharmacology*, 13, 1008580.
17. Yin, X., et al. (2024). Dietary factors influence gut microbiota composition and aging pathways in *Drosophila melanogaster*. *Frontiers in Microbiology*.
18. Huang, J. H., et al. (2015). Role of gut bacteria in regulating lipid metabolism under high-sugar diet in *Drosophila*. *Royal Society Biology Letters*.
19. Gálíková, M., & Klepsatel, P. (2018). Obesity and aging in the *Drosophila* model. *International Journal of Molecular Sciences*, 19(7), 1896. <https://doi.org/10.3390/ijms19071896>